

Demonstration 2(em):Einstein-Maxwell equations in the Kerr-Newman spacetime.

> **restart:**

> **grtw():**

GRTensorII Version 1.79 (R6)

2 February 2001

Developed by Peter Musgrave, Denis Pollney and Kayll Lake

Copyright 1994-2001 by the authors.

Latest version available from: <http://grtensor.phy.queensu.ca/>

> **grOptionTermSize:=30:**

> **qload(newkn);**

Default spacetime = newkn

For the newkn spacetime:

Coordinates

x(up)

$x^a = [r, u, \phi, t]$

Line element

$$ds^2 = \frac{(r^2 + u^2) dr^2}{r^2 - 2mr + a^2 + Q^2} + \frac{(r^2 + u^2) du^2}{a^2 - u^2} + \frac{(a^2 - u^2) \left(r^2 + a^2 + \frac{(a^2 - u^2)(2mr - Q^2)}{r^2 + u^2} \right) d\phi^2}{a^2} - \frac{2(a^2 - u^2)(2mr - Q^2) d\phi dt}{a(r^2 + u^2)} + \left(-1 + \frac{2mr - Q^2}{r^2 + u^2} \right) dt^2$$

Constraints = [u = a cos(theta)]

*Kerr Newman Solution in Boyer-Lindquist coordinates (u=a*cos(theta))*

a)Background:

Define the vector potential \$A_{[a]}\$ and the electromagnetic field tensor \$f_{[ab]}\$:

> **grdef(`A{a}:=[0,0,Q*r*(a^2-u^2)/(a*(r^2+u^2)), -Q*r/(r^2+u^2)]`);**

Components assigned for metric: newkn

Created definition for A(dn)

> **grdisplay(A(dn));**

For the newkn spacetime:

A(dn)

A(dn)

$$A_a = \left[0, 0, \frac{Qr(a^2 - u^2)}{a(r^2 + u^2)}, -\frac{Qr}{r^2 + u^2} \right]$$

> **grdef(`f{[a b]}:=2*A{[b ; a]}`);**

Created a definition for A(dn,cdn)

```

Created definition for f(dn,dn)
Two scalar invariants associated with the field tensor are  $f_{[ab]}f^{[ab]}$  and  $f_{[ab]}f^{*ab}$ :
> grdef(`I1:=f{a b}*f{^a^b}`);
Created definition for f(up,up)
Created definition for I1
> grdef(`fstar{a b}:=LevC{a b c d}*f{^c^d}`);
Created definition for fstar(dn,dn)
> grdef(`I2:=f{a b}*fstar{^a^b}`);
Created definition for fstar(up,up)
Created definition for I2
The 4-current is defined by  $J^a=f^{ab}_{;b}/(4\pi)$ 
> grdef(`J{^a}:=f{^a^b;b}/(4*Pi)`);
Created a definition for f(up,up,cdn)
Created definition for J(up)
and for the field tensor we construct the permutation  $f_{[ab;c]}+f_{[bc;a]}+f_{[ca;b]}$ .
> grdef(`Id{a b c}:=f{a b ; c}+f{b c ; a}+f{c a ; b}`);
Created a definition for f(dn,dn,cdn)
Created definition for Id(dn,dn,dn)
b)Calculations: First we calculate, simplify and display the field tensor and invariants. The four-current and permutaion of the field tensor are then reduced to zero.
> grcalc(f(dn,dn));

CPU Time = .080

> gralter(_,2,7);
Component simplification of a GRTensorII object:

Applying routine `simplify[trig]` to object f(dn,dn)
Applying routine factor to object f(dn,dn)

CPU Time = .030

> grdisplay(_);

For the newkn spacetime:


$$f(dn,dn)$$


$$f(dn, dn)$$


$$f_{r\phi} = -\frac{(r-u)(r+u)Q(a-u)(a+u)}{a(r^2+u^2)^2}$$


$$f_{rt} = \frac{Q(r-u)(r+u)}{(r^2+u^2)^2}$$


$$f_{u\phi} = -2\frac{Qru(a^2+r^2)}{a(r^2+u^2)^2}$$


$$f_{ut} = 2\frac{Qru}{(r^2+u^2)^2}$$


> grcalc(I1,I2);

CPU Time = .060

```

```
> gralter(_,2,7,10);
```

Component simplification of a GRTensorII object:

Applying routine `simplify[trig]` to object I1

Applying routine `simplify[trig]` to object I2

Applying routine factor to object I1

Applying routine factor to object I2

Applying routine `simplify[sqrt]` to object I1

Applying routine `simplify[sqrt]` to object I2

CPU Time = .060

```
> grdisplay(_);
```

For the newkn spacetime:

I1

$$I1 = -2 \frac{Q^2 (-u^2 + 2 r u + r^2) (-u^2 - 2 r u + r^2)}{(r^2 + u^2)^4}$$

I2

$$I2 = 16 \frac{(r - u) (r + u) Q^2 r u}{(r^2 + u^2)^4}$$

```
> grcalc(J(up));
```

CPU Time = .050

```
> grdisplay(_);
```

For the newkn spacetime:

J(up)

J(up)

J^a = All components are zero

```
> grcalc(Id(dn,dn,dn));
```

CPU Time = .070

```
> grdisplay(_);
```

For the newkn spacetime:

Id(dn,dn,dn)

Id(dn,dn,dn) = All components are zero

c)Einstein-Maxwell equations: We construct the energy-momentum tensor out of the field tensor ($T_{ab} = (f_{ac}f_b^c - g_{ab}f_{cd}f^{cd})/4 - 8\pi T_{ab}$) and then show that the **Kerr-Newman spacetime** satisfies the Einstein-Maxwell equations.

```
> grdef(`T{a b} := (+f{a c}*f{b^c} - g{a b}*I1/4)/(4*Pi)`);
```

Created definition for f(dn,up)

Created definition for T(dn,dn)

```
> grdef(`Dif{a b} := G{a b} - 8*Pi*T{a b}`);
```

Created definition for Dif(dn,dn)

```
> grcalc(Dif(dn,dn));
```

CPU Time = .090

```
> gralter(_,6,7);  
Component simplification of a GRTensorII object:  
  
Applying routine expand to object Dif(dn,dn)  
Applying routine factor to object Dif(dn,dn)  
  
CPU Time = .010  
  
> grdisplay(_);  
  
For the newkn spacetime:  
Dif(dn,dn)  
Dif(dn, dn)  
Difa b = All components are zero  
  
>
```